

## Stratification and disconformities in yellow sands of the Bassendean and Spearwood Dunes, Swan Coastal Plain, South-western Australia

D K Glassford<sup>1</sup> & V Semeniuk<sup>2</sup>

<sup>1</sup>33 Rockett Way, Bull Creek, WA 6155

<sup>2</sup>21 Glenmere Road, Warwick, WA 6024

Manuscript received September 1988; accepted January 1989

### Abstract

Yellow sand of the Pleistocene Bassendean and Spearwood dunes of the Swan Coastal Plain of SW Australia contains sedimentary stratification at depth (>3-5m). Yellow sand is divided into units of high and low angled inclined strata, horizontal strata, and massive sand. Sequences of sand typically contain up to five major disconformities. Dip direction resultants of inclined strata at individual study sites indicate that Pleistocene wind directions were dominantly from the eastern sector, sub-dominantly from the NW, and to a minor extent from the W and SW. Dip direction mean vector resultant of all inclined strata is 210° indicating that the sands were transported mainly from the NE. These data preclude, as major processes, origin of the yellow sand by either *in situ* decalcification of coastal limestones or by coastal-marine derivation as carbonate-free quartz sand. Cross-strata and horizontal strata were emplaced by the migration of desert aeolian dune fields (desert ergs) during at least six glacial age arid phases, mostly during the middle Pleistocene. During the intervening wetter interglacial periods yellow sands were extensively bioturbated to depths of 3-5m, thereby degrading the dunes and producing massive yellow sand under the major disconformities.

### Introduction


The Swan Coastal Plain of the Perth Basin, South-western Australia has an extensive cover of mainly Pleistocene siliciclastic sand (McArthur & Bettenay 1960, Playford *et al* 1975, 1976; Wilde & Low 1978, 1980; Fig. 1). This cover includes Bassendean Sand, yellow sand overlying Tamala Limestone, and some of their altered equivalents such as white quartz sand, brown quartz sand and brown sandstone. These sand formations traditionally have been interpreted as coastal-marine derived calcareous deposits which have been decalcified *in situ* by leaching processes to a residue of yellow quartz sand (Prider 1948, McArthur & Bettenay 1960, Lowry 1977, Wyrwoll & King 1984, amongst others).

Crucial evidence of long standing for decalcification (*eg* Prider 1948) is, that near the coast where yellow sand overlies limestone, from which traditionally it is concluded the sand was leached, the yellow sand has a massive structure (*eg* Spearwood Dunes). Inland where yellow sand does not overlie limestone it generally also has a similar massive structure (*eg* Bassendean Sand, yellow sand of the Yoganup Formation). Thus massive structure (absence of lamination/ bedding) has been cited as key evidence for an *in situ* decalcification origin. However, documentation of these sand formations in terms of their genetically important sedimentary features (*eg* stratigraphy, geometry, structure, fabric, texture and composition) is negligible.

This paper provides the first detailed treatment of primary sedimentary structures in the yellow sands of the Bassendean and Spearwood dunes within the central sector of the Swan Coastal Plain (Fig. 1). The paper deals only with Bassendean Sand and the quartz sand overlying the Tamala Limestone referred to the Karrakatta and Cottesloe soil associations (McArthur & Bettenay 1960), and specifically excludes the Cooloongup Sand with its local grey coloration and occasional shell content (Passmore 1970), yellow sand of the Yoganup Formation (Low 1971), Eaton Sand (Semeniuk 1983), the complex quartz sand deposits of the hinterland of Geographe Bay (Baxter 1977), and possible minor non-aeolian facies of the yellow sands (*eg* Wyrwoll & King 1984). It is also stressed that this study mainly concentrates on primary sedimentary features of the yellow sands and not on their pedogenic or geohydrologic alteration products such as humic quartz sand, white quartz sand and ferruginized sand. These various secondary overprints are briefly described as a background to understanding the destruction of primary sedimentary structures in yellow sands.

### Previous studies

The major geological attributes of Perth Basin yellow sand have not been documented in systematic detail. These attributes should include genetically critical sedimentary features such as geomorphic expression, boundary types, stratigraphic associations and relationships, ge-



**Figure 1** Location and regional setting of study sites. **A** Location of study area in Western Australia. **B** Location of study area in the Perth Basin. **C** Regional geomorphic units of the Swan Coastal Plain, the landward portion of the central Perth Basin, and their relationship to insets of study site settings (after McArthur and Bettenay 1960). **D, E, F & G** Location and setting of study sites (geomorphic unit boundaries after Gozzard 1983, 1986; Jordan 1986). Note the boundary between the Bassendean and Spearwood Dune sands as mapped by Gozzard (1983, 1986) and Jordan (1986) is 1.5 to 4km further east than it is as mapped by McArthur & Bettenay (1960).

ometry and dimensions, internal structure (types, dip angles, dip directions *etc*), fabric, texture (grain size, sorting, percentage fines, roundness, *etc*) and composition (grain types, grain coatings, internal features of grains, mineralogy of grains *etc*) (eg see Folk 1974, Pettijohn *et al* 1987, Lindholm 1987). Previous investigations into yellow sand have been directed towards agriculture and pedology (McArthur & Bettenay 1960), environmental and land use planning (Dept of Conserv & Envir 1980), hydrology and groundwater supply (Allen 1981), engineering (Klenowski 1976), heavy mineral deposits (Baxter 1977), geological mapping (Wilde & Low 1978, 1980; Gozzard 1983, 1986; Jordan 1986); and lithostratigraphic sub-division and definition (Low 1971, Playford & Low 1972). These studies provide information on various aspects of yellow sand, but they are not directly relevant to unravelling its origin. Despite this, many of these studies form the basis for the traditional views of coastal-marine derivation and *in situ* decalcification for the origin of yellow sand (Lowry 1977). Two published studies have dealt specifically with the genesis of Perth Basin yellow sand (Glassford & Killigrew 1976 and Wyrwoll & King 1984). These studies, however, focused mostly on the textures of yellow sand.

Although stratification is an important feature for interpreting the origin of sand (Potter & Pettijohn 1977, Lindholm 1987), it has not been the subject of any previous work on Perth Basin yellow sand. Generally, yellow sands are regarded to be structureless or massive (eg Prider 1948, Wyrwoll & King 1984). The only published mention, to our knowledge, of stratification in yellow sand is the reference by Baxter (1977) to "delicate cross-bedding" in Bassendean Sand in the Busselton area.

### Methods

Stratification and unconformities in yellow sand sequences were studied in 15 quarry exposures in the Perth region of the Swan Coastal Plain (Fig. 1). These quarries represent all sand quarries that to our knowledge were being excavated during this study. Stratification was best studied in winter when the sand is moist. In summer, the sand is dry and commonly slumps during and following excavation. This slumping covers the deeper sequences of yellow sand, thereby typically concealing stratification. Four of the quarries are located in the middle to eastern parts of the Spearwood Dunes; eleven of the quarries are in the Bassendean Dunes. Structures within the sand were documented using the terminology of McKee & Weir (1953) and Lindholm (1987). Stratification and unconformities were mapped in the field onto panoramic photographs of quarry walls.

The true dip angle and dip direction of cross-layering was determined by measurements on partially excavated individual surfaces of the sedimentary layers or laminae of a given cross-layered set which were traceable in three dimensions. Usually 5-10 separate laminae were measured in this manner at each sampling locality. Depending on the amount of exposure, 2-5 sites typically were measured in each quarry. Dip angle and dip direction (magnetic) readings were taken according to the number and thickness of large scale cross-layered sets. Most readings were at c 0.5-1.5m vertical intervals for a set. For example, for a 3m thick set, three readings were taken at approximately 0.5m, 1.5m and 2.5m above the base of the set. In the case of a 5m thick set five readings were taken at approximately, 0.5m, 1.5m, 2.5m, 3.5m and 4.5m above the base of the set. The constraints of availability and wall condition of quarries has limited the data base of this paper to 228 dip angle and direction measurements of inclined strata from 12-15 lo-

calities. Additional measurements should be collected as existing quarries are excavated further and when new quarries are opened.

Sketches and photographs were used to record secondary overprints. Sediment was carefully sampled *in situ* by a 5 cm corer and retrieved to the laboratory for impregnation and slicing. Oriented thin sections were prepared from the impregnated *in situ* blocks for petrographic study. Selected laminae and groups of laminae also were sampled and sieved at half phi intervals for granulometric analysis. Representative samples and sub-samples were x-rayed with Co K-alpha radiation and diffractograms were interpreted using Brindley & Brown (1980).

### Nomenclature

The Sahara Desert term *erg* is used in this study. Originally *erg* referred to a vast region in the Sahara Desert covered by deep aeolian sand (Gary *et al* 1972). Implicit in this original meaning were a mid-latitude location, a desert setting, and a continental, as distinct from coastal, character and derivation for the sand. Subsequent workers, however, expanded the original meaning of the term to encompass large scale coastal and non-desert aeolian sand bodies, or sand seas (eg Blakey *et al* 1988; Marzolf 1988). It is therefore important to define the sense in which *erg* is used. In this study *erg* refers to a vast, large scale or regional scale tract of continental desert-aeolian sand in the form of sheets and dunes. For added emphasis and clarity the term is used in conjunction with *desert*, ie *desert erg*. Other types of aeolian sand seas can then be categorized, for example, as *coastal erg* or *humid erg*.

The term *desert* is used in this paper in the sense of low to middle latitude terrains with no vegetation cover or with a vegetation cover which is too sparse to prevent widespread aeolian transport of sand. Controls such as effective rainfall, windiness, sand supply *etc* may variously contribute to producing a desert terrain. For a discussion of the main factors which can produce a desert-aeolian terrain see Ash & Wasson (1983) and Marzolf (1988).

### Geological setting

The study area is within the Swan Coastal Plain, a coastal lowland of the Perth Basin with a mainly relict Pleistocene surface (Playford *et al* 1976, Wilde & Low 1978, 1980). Within this setting there is a Pleistocene lithostratigraphic unit of yellow sand (Prider 1948, Glassford 1980) relevant to this study which usually has been assigned to more than one formation (Fig. 1):

- yellow quartz sand portion of Bassendean Sand (= Bassendean Dunes of McArthur & Bettenay, 1960)
- yellow quartz sand assigned to the Tamala Limestone (within the Spearwood Dunes of McArthur & Bettenay, 1960).

To date, however, the bulk of the yellow quartz sand overlying the Tamala Limestone within the Spearwood Dunes has not been formally recognized as a separate formation, except in local areas such as Rockingham (Coo loongup Sand; Passmore 1970) and Australind (Eaton Sand; Semeniuk 1983). Lithologically similar yellow sands (Prider 1948, Glassford 1980) which are located east of Perth on the eastern margin of the Pinjarra Plain and in the foothills of the Darling Scarp (Low & Lake 1970, Low *et al* 1970, Wilde & Low 1978, 1980) have been assigned to the Yoganup Formation (Low 1971). For the sake of simplicity the Yoganup Formation yellow sands in this complex geomorphic setting are excluded from the present study.

The age of the yellow sands is interpreted to be mainly middle Pleistocene (Playford *et al* 1975). It forms much of the land surface of the coastal plain to landward of the Holocene coastal dune belt and the Pleistocene limestone ridges. Where the yellow sand underlies the coastal limestones it may range to Pliocene in age (Logan *et al* 1970, Glassford 1980).

The terrain of yellow sand on the coastal plain is the western fringe of a vast, diachronous cover of sand dunes and sand sheets which extend westwards from the central Australian Desert (Killigrew & Glassford 1976, Glassford 1980, 1987, Beard 1984). Thus yellow sand occurs extensively on the Yilgarn Block and forms an extensive and thick lithosome that underlies and overlies coastal limestones and underlies landward parts of the coastal plain (Semeniuk & Glassford 1988).

On the coastal plain near the present coast, there is a shore-parallel belt of Pleistocene coastal limestone ridges (Tamala Limestone). These ridges, with variable cover of yellow sand, form shore outcrops, or may be buried by Holocene littoral and coastal-dune deposits. In cross section the coastal limestone complex forms a large lens mainly on, but also within, the western margin of the aeolian sand sheet described above, a relationship that is evident for much of the west coastal region of Western Australia from Shark Bay to Perth, a distance of over 1 200 km (Logan *et al* 1970, McWhae in Quilty 1974, Glassford 1980, Allen 1981, Semeniuk & Glassford 1988). The eastern contact of the limestone-ridge-belt with yellow quartz sand of the hinterland is complicated: it may be sharp, or lithologically gradational, marked by east-west inter-fingering, or marked by zones of scattered limestone lenses in yellow sand (Semeniuk & Glassford 1988).

The lithofacies referred to herein as yellow sand, although typically yellow, also includes sands varying from red to brown to locally white. Grain sizes range from bimodal to poorly unimodal to unimodal, and the sand is typically fine skewed, well to poorly sorted and medium sized. Framework grains are predominantly quartz, with moderate to minor microcline and minor heavy minerals. The grains have a coating of silt-clay sized kaolin, goethite (yellow) and/or haematite (red) and quartz. These latter minerals also occur in the less than 0.09mm fraction of the sediment (Glassford 1980).

#### Description and interpretation of stratification in yellow sand

Contour data from 1:25 000 topographic sheets and field observations of surface morphology indicate that sites 1, 2, 3, 4 and 7 of this study are within large scale sand-sheet ramparts which drape the landward side of the coastal-limestone-ridge belt; sites 5 & 6 are in sinuous and parabolic landforms; and sites 8 to 15 are within an east-west oriented linear dune with stellate landforms (Table 1; Fig. 1). The term *draa* refers to large scale dunes or megadunes and compound or complex types of composite dunes (Wilson 1972, Leeder 1982, Lancaster 1988).

On the basis of presence/absence and inclination of large scale (> 1m thick) sets of stratification, yellow sand is divisible into:

- 1) units with inclined strata ie cross-strata (high and low angled);
- 2) units with horizontal strata; and
- 3) units with massive structure.

These features are described below. Small scale (<0.3m thick) sets of cross-strata were not observed at any of the sites. The thinnest recognizable strata or laminae that comprise sets of cross-strata and horizontal strata in yellow sand are mostly evident due to differences in size of framework sand grains and, to a lesser extent, to varying amounts of yellow to brown silt-clay sized matrix fill of kaolin and goethite and variations in framework-grain coatings of kaolin and goethite.

#### Inclined strata

##### Description

Large scale sets of inclined strata (commonly referred to as cross-strata) occur extensively in 13 of the quarries (Table 1). Sets of cross-strata typically are up to 5-10m thick, and generally, only occur at depths of more than 3-5m below the present land surface. Cross-stratification consists dominantly of tabular-shaped sets of planar cross-strata with local minor occurrences of wedge-shaped and hummocky-shaped sets of planar and curved cross-strata (Figs 2, 3 & 4). Tabular-shaped sets of planar cross-strata have been traced continuously along recently exposed quarry faces for distances of 10 to 100 m. Generally exposures of cross-stratification are limited by an upper zone of bioturbation, or by lateral slumps of quarry walls, or by quarry floors.

Table 1



#### Sand quarries near Perth with large scale cross-stratification and major disconformities.

Quarries examined and their site numbers <sup>1</sup>	Large scale cross-stratification <sup>2</sup>	Major disconformities <sup>3</sup>
1 Wangara North	present	2 present
2 Wangara	present	3 present
3 Landsdale West	-	2 present
4 Landsdale East	-	1 present
5 Belhus	present	-
6 Henley Brook	present	1 present
7 Mirrabooka	present	5 present
8 Jandakot West	present	-
9 Jandakot East	present	-
10 Jandakot North 1	present	-
11 Jandakot North 2	present	-
12 Canning Vale	present	1 present
13 Banjup West	present	1 present
14 Banjup East 1	present	1 present
15 Banjup East 2	present	1 present


<sup>1</sup> See Fig. 1 for location of sites.

<sup>2</sup> Large-scale is defined as greater than 1.0m thick.

<sup>3</sup> Number of major disconformities in the yellow sand sequences.




**Figure 2** Range of stratification features at a single typical locality, Banjup West. See Fig. 1 for location. **A** Panoramic diagram of stratification and location of insets. **B** A large scale, tabular set of planar horizontal strata overlain by a large scale, wedge-shaped set of planar cross-strata (inclined strata). **C** A large scale set of concave-upwards, trough cross-strata. Note the rivulets of flowing sand which are building avalanche cones of sand. The continued growth of avalanche cones of sand will eventually conceal the stratification. **D** Thinnest recognizable strata in a large-scale set of low angle, planar cross-strata produced by differences in grain size of framework sand grains. The coarser layers (a) resemble ripple-form strata of Hunter (1977). The finer layers (b) resemble grainfall strata of Hunter (1977). Coin is 21mm in diameter. **E** Layering in a large-scale set of planar cross-strata produced by differences in texture, composition and colour. Thin prominent layers have framework quartz-sand grains supported in a silt-clay matrix of yellowish brown goethite, kaolin and quartz. Thick layers lack the silt-clay matrix. Coin is 28mm in diameter. **F** Discontinuity (a) between two units of yellow sand. The discontinuity between the units is produced by: geometry of interface; changes in stratification details away from the contact; and structural, textural, compositional and colour differences between the two units on either side of the interface. The massive zone below the discontinuity is a palaeosol (b) which indicates the discontinuity is a regional bounding surface (Talbot 1985). Note also the common indistinct appearance of stratification. Several metres of surficial sand has been removed from this section by excavating machinery. Spade for scale is 1 m in length. **G** Closer view of disconformity surface, sloping from upper left to lower right and with underlying, mostly massive palaeosol with rare organic material and burrow-structures (a). Note rivulets of flowing sand on the left hand side. Coin is 24 mm in diameter.



**Figure 3** Panoramic diagrams of stratification attitudes and relationships within sequences of yellow sand, Swan Coastal Plain, southwestern Australia. See Fig. 1 for locations.



**Figure 4** Primary stratification and secondary overprint features in yellow sand. See Fig. 1 for locations. **A** Four units (1, 2, 3, 4) of yellow sand bounded by three unconformities (I, II, III), Mirrabooka quarry. Note the common indistinct nature of the stratification. Unit 1 is mostly massive. Unit 3 has large scale, high angle planar and tangential cross-strata which dip towards the west (towards RHS). Unit 3 has large scale, high and low angle planar cross-strata which dip towards the east (towards LHS). Unit 4 has been truncated by earth-moving machinery. Spade for scale is 1 m in length. See Table 2 & Fig. 7. **B** Set of large scale, low angle, planar cross-strata dipping towards the southwest (towards LHS), Jandakot North 2. Note firstly, cross-stratification at depth is obscured by cone-shaped slumps of sand along the base of the cross-stratified sand escarpment. Secondly presence of an overlying soil of white and black sand which has been disrupted and partly removed by earthmoving machinery. Set is 7-8 m thick. **C** Set of large scale, broadly concave-upwards, trough cross-strata, Jandakot West. Note 28 mm diameter coin in lower left hand corner, below a broad trough-shaped bounding surface (a). Stratification is obscured by slumped sand on the lower right hand side. **D** Large-scale set of low angle and horizontal strata which resemble grainfall and plane bed strata of Hunter (1977). Strata underlie an interdune area, Canning Vale. Coin 28mm in diameter. **E** Rare cut-and-fill structure, Banjup West. Lower set of horizontal strata has been truncated by erosion (? surface wash) and then overlain by an upper westward dipping set of large-scale high angle, tangential to planar cross-strata. **F** Burrow structure (a) in horizontally stratified sand which has negligible goethite and kaolin silt-clay, Jandakot North. Layers resemble grainfall and plane bed strata of Hunter (1977). Coin is 28mm in diameter. **G** Termitarium structured yellow sand, upper few metres of unit 4, Mirrabooka. In contrast to F above the sand is relatively rich in goethite and kaolin silt-clay (see Table 2). **H** Large-scale set of low angle, planar cross-strata (lower one third) overlain by a large-scale set of horizontal strata (upper two thirds) which underlie an interdune and are partly ferruginized to form brown sandstone (*coffee rock*).



**Figure 5** Frequency distribution of true dip angles for high angle (equal or greater than  $20^\circ$ ) and low angle ( $1.6$ - $19^\circ$ ) cross-strata and horizontal strata ( $0$ - $1.5^\circ$ ) from yellow sands of the Perth area, Swan Coastal Plain, southwestern Australia.

Dip angles of cross-strata have a bimodal frequency. High angle (equal to or greater than  $20^\circ$ ) cross-strata comprise c 51% of measurements and form a mode between  $25$ - $30^\circ$  (Fig. 5). The highest dip angle recorded for cross-strata in yellow sand is  $35^\circ$ . Low angle ( $1.6$ - $19^\circ$ ) cross-strata comprise 49% of measurements and form a mode between  $5$ - $10^\circ$ , and the mode is skewed towards  $10$ - $15^\circ$  (Fig. 5). The proportion of high angle strata is probably lower than these figures indicate because of a bias in the location of quarries towards hill tops rather than plinths and interdune flats.

The frequency of occurrence of dip directions (azimuths) of cross-strata at individual sites may be unimodal, bimodal or polymodal (Fig. 6). Individual disconformity-bounded sets of cross-strata at the same locality may have variable dip directions (Table 2; Fig. 7). At six sites cross-strata have mean resultant dip direction towards the W and SW. At three sites cross-strata have mean resultant direction towards the E and SE. At two sites cross-strata have mean resultant dip direction towards the NE. At one site the resultant dip direction changes between the various disconformity bounded units of yellow sand (Fig. 6). Dip



Table 2


Description of lithology for the sequence of yellow sands at Mirrabooka quarry. See Fig. 1 for location. Grain size terminology after Folk (1974).

DEPTH (m)	DESCRIPTION	LITHOFACIES
0 to 4.0*	Quartz sand; yellow, massive, sand-framework supported, unimodal, near symmetrical to fine skewed (range +0.0 to +0.12), well sorted to moderately well-sorted (range 0.47 to 0.61 phi), medium (range 1.21 to 1.63 phi) quartz sand with 0.9% fines** (range 0.60 to 1.18%); framework quartz-sand grains of yellow sand have a surface coating of goethite (yellow) pigmented kaolin plus silt-clay sized quartz	Unit 6
DISCONFORMITY (V)**		
4 to 9	Quartz sand; white in upper 0 to 0.3m then yellow to 9m, massive, sand-framework supported, unimodal, near symmetrical (range +0.02 to -0.05), moderately well-sorted (range 0.58 phi to 0.60 phi), medium (range 1.56 to 1.63 phi) quartz sand with 1.18% fines (range 0.97 to 1.35%); framework quartz-sand grains of yellow sand have a surface coating of goethite (yellow) pigmented kaolin plus silt-clay sized quartz; the white sand does not have a yellow coating on grains.	Unit 5
DISCONFORMITY (IV)		
9 to 14	Quartz sand; in places upper 0 to 0.3m white then yellow and yellowish red with white mottles to 14m; in places upper 0 to 0.5-1.0m is termitarium structured with a tortuous network of vermiform voids or loose sand fill and labyrinthoid walls of coherent sand and silt-clayey sand; from 9-10 to 12.5m conspicuously cross-stratified with large scale high angle strata dipping towards the southeast; from 12.5 to 14m conspicuously cross-stratified with large scale low angle strata dipping towards the southwest; sand-framework supported, unimodal to poorly unimodal, fine skewed to near symmetrical (range +0.15 to -0.05), moderately well-sorted to moderately sorted (range 0.54 to 0.72 phi), medium (1.33 to 1.66 phi) quartz sand with 1.55% fines (range 0.88 to 2.08%); in places thin beds of coarse sand, framework quartz-sand grains of yellow to red sand have a surface coating of goethite, haematite, kaolin and silt-clay sized quartz; white sand does not have coating on grain surfaces.	Unit 4
DISCONFORMITY (III)		
14 to 17.5	Quartz sand, brownish yellow to reddish yellow to brown with white mottles; conspicuously cross-stratified, from 14 to 17.5m large scale, high and low angle strata dip towards the northeast and east, in places from 16 to 17.5m large scale, low angle cross-strata dip towards SSE; sand-framework supported, unimodal, near symmetrical (range +0.03 to -0.06), moderately well-sorted (range 0.52 to 0.60 phi), fine to medium (range 1.69 to 2.03 phi) quartz sand with 2.24% fines (range 1.84 to 2.74%); framework quartz-sand grains of yellow to brown sand have a surface coating of goethite and kaolin plus silt-clay sized quartz; white sand mottles have grains which do not have surface coatings	Unit 3
DISCONFORMITY (II)		
17.5 to 20.5	Quartz sand; pale yellow to reddish yellow; generally massive, in places faintly cross-stratified with high angle strata dipping towards the W & WNW; sand-framework supported, unimodal, near symmetrical to strongly fine-skewed (range +0.08 to +0.61), moderately well-sorted (range 0.64 to 0.69 phi), medium (range 1.41 to 1.7 phi) quartz sand with 2.85% fines (range 2.53 to 3.13%); framework quartz-sand grains of yellow sand have a surface coating of goethite, haematite, kaolin and silt-clay sized quartz	Unit 2
DISCONFORMITY (I)		
20.5 to 26+	Quartz sand; yellow to red, white in the vicinity of the ground water table (c 24-26m); generally massive, in places faintly cross-stratified with large scale, high angle strata dipping towards the NW, sand-framework supported, unimodal to poorly unimodal, near symmetrical to strongly fine skewed (range +0.06 to 0.37), moderately well sorted to moderately sorted (range 0.61 to 0.97 phi), coarse to medium (range 0.97 to 1.36 phi) quartz sand with 3.15% fines (range 1.12 to 6.57%); framework quartz-sand grains of yellow to red sand have a surface coating of goethite (yellow) or haematite (red) pigmented kaolin plus silt-clay sized quartz, white sand in vicinity of groundwater table has grains which do not have a surface coating; groundwater table c 0.5m below base of quarry.	Unit 1


\* Natural surface has been removed

\*\* Fines equal weight percentage of grain size less than 0.09mm. Generally fines of yellow to red sand are composed of c 50% kaolin, goethite and/or haematite and c 50% quartz and heavy minerals. Fines of white sand are composed of quartz and heavy minerals. Note that there is a uniformly consistent decrease in the amount of fines from unit 1 at depth to unit 6 near the surface. Furthermore modern beach-berm sands average 0.09% fines (range 0-0.48%) and Holocene coastal dunes average 0.15% fines (range 0-0.62%), see details in Glassford (1980)

\*\*\* Discontinuities recognised by occurrence of palaeosols, undulating discontinuity surfaces, changes in structure (eg massive to cross-stratified) and changes in fabric, texture, composition and colour.



**Figure 6** Rose diagrams of dip directions of cross-strata for individual sites in the Perth area, Swan Coastal Plain, southwestern Australia. See Fig. 1 for locations. Map boundaries are after Wilde and Low (1978, 1980).



**Figure 7** Vertical section of six disconformity-bounded units of yellow sand and the dip angles and dip directions of cross-strata for the respective lithostratigraphic units of yellow sand. Note that the yellow sands are massive near the surface stratified at depth and then massive near the groundwater table.

directions may vary markedly between and within units of yellow sand at the one site. For instance at Mirrabooka (Fig. 1) the lower strata of unit 4 dip towards 240° whereas middle to upper strata dip towards 160° (Table 2; Figs 6 & 7).

The dip directions of all cross-strata have a polymodal frequency distribution. High angle and low angle cross-strata also have a polymodal frequency distribution (Fig. 8). High angle cross-strata have prominent modes between 135° and 171°, between 243° and 261°, skewed to 279° and between 297° and 315°, skewed to 279° (Fig 8A). Low angle cross-strata have prominent modes between 135° and 153°, skewed to 171°, between 189° and 207°, skewed to 225° and between 297° and 315° (Fig. 8B).

The mean vector resultant is (Figs 8 & 9):

- towards 220° for high angle, large scale cross-strata
- towards 198° for low angle, large scale cross-strata and
- towards 210° for all large scale cross-strata.

*Interpretation*


High angle, large scale cross-strata are interpreted to be the lee-slope foresets of aeolian dunes (following McKee & Bigarella 1979, Nielson & Kocurek 1987). Dip directions of the high angle cross-strata indicate deposition by SE winds, ENE winds and NNW winds (Table 3). Low angle cross-strata are interpreted to be predominantly a combination of lee-slope foresets, dune (including star dune) foot-slope plinths and sand-sheet sets (following Ahlbrandt & Fryberger 1981, Kocurek 1981, Nielson & Kocurek 1987). Dip directions are roughly similar to those of high angle cross-strata and indicate deposition by SE winds, NNE winds, and NW winds (Table 4). Mean vector resultants for high angle, low angle, and high angle plus low angle cross-strata indicate long-term overall sand movement, on average, was from NE towards SW (Table 5; Figs 8 & 9). This is opposite to the direction of transport of carbonate sands of the present modern coastal dunes, which is towards the NE (Searle & Semeniuk 1985).

**Table 3**


**Formative wind directions and predominant direction of sand movement for large scale high angle cross-strata.**

Modal dip directions of large scale high angle cross-strata <sup>1</sup>	Formative wind direction	Direction of sand movement
Towards 306° (towards NW)	From SE	From SE to NW
Towards 252° (towards WSW)	From ENE	From ENE to WSW
Towards 162°-144° (towards SSE)	From NNW	From NNW to SSE

<sup>1</sup> Modes are from Fig. 8



**Figure 8** Dip direction plots of cross-strata in yellow sands of the Perth area, Swan Coastal Plain, southwestern Australia. **A** Rose diagram of dip directions of high angle (equal to or greater than 20°) cross-strata. **B** Rose diagram of dip directions of low angle (1.6-19°) cross-strata. **C** Scatter plot diagram of dip angles and dip directions for all cross-strata. **D** Rose diagram of dip angles and dip directions for all cross-strata. Note **A** & **B** are plotted with an arithmetic scale, and not an equal-area scale (cf Nemec 1988).



**Figure 9** Graph of vector magnitude (consistency ratio of Reiche 1938) versus number of observations of dip direction to show significance (at 0.05) of the vector magnitude (following Lindholm 1987). Note the vector magnitudes which plot above the 0.05 probability line, indicate there is less than 1 chance in 100 that the population does not have a preferred direction of movement.

**Table 4**

Formative wind directions and predominant direction of sand movement for large scale low angle cross-strata.

Modal dip directions of large scale low angle cross-strata <sup>1</sup>	Formative wind direction	Direction of sand movement
Towards 306° (towards NW)	From SE	From SE to NW
Towards 216°-199° (towards SSW)	From NNE	From NNE to SSW
Towards 144° (towards SE)	From NW	From NW to SE

<sup>1</sup> Modes are from Fig. 8

**Table 5**

Directions of mean vector resultants for high angle, low angle, and high angle plus low angle cross-strata and their palaeo-directional interpretations.

Cross-strata	Direction of mean vector resultant <sup>1</sup>	Interpretation
High angle	Towards 220° (towards SW)	Overall sand movement is from NE to SW under the influence of Pleistocene winds which were predominantly from the eastern sector
Low angle	Towards 198° (towards SSW)	Overall sand movement is from NNE to SSW under the influence of Pleistocene winds which were predominantly from the eastern sector
High angle plus low angle	Towards 210° (towards SSW)	Overall sand movement is from NNE to SSW under the influence of Pleistocene winds which were predominantly from the eastern sector

<sup>1</sup> Vector resultants are from Fig. 8.

The overall sand movement from NE has many local variations (Figs 6 & 7). Sites which are closely located (eg 10, 11, 12, 13 & 14; Fig. 6) may have a wide divergence in direction of their dip direction resultants. Furthermore the dip directions of individual sites (eg sites 8-14; Fig. 6) are also widely divergent, typically over more than 180°. Interconnected (by one or more arms) stellate dune morphology, plus the associated wide range in dip directions of the internal strata of the same features, suggests that sites 8-14 are located within a linear draa (or megadune) with star dunes. This interpretation is important, because the internal structures of star dunes are imperfectly known and only rarely have they been identified in the sedimentary record (Nielsen & Kocurek 1987).

It has been previously suggested elsewhere that yellow sand is predominantly glacial age desert-aeolian sediment (Glassford & Killigrew 1976, Killigrew & Glassford 1976, Glassford 1980, Semeniuk & Glassford 1988). Cross-strata details (Figs 6, 7, 8 & 9) provide additional support for a desert-aeolian origin and are consistent with the following refinements to the desert aeolian model in terms of more specific formative wind directions. During periods of deposition of yellow sands, Pleistocene wind directions were dominantly from the eastern sector, sub-dominantly from the NW and to a minor extent from the W and SW.

### Horizontal strata

#### Description

Horizontal strata (c 0-1.5° angle of dip) comprise c 13% of all strata measured (Fig. 5). As discussed later this low proportionate amount is probably an artifact of the location of quarries in hills of sand rather than sand flats. Sets of horizontal strata are up to 7 m thick, and have been traced laterally for up to 50 m and, by interpolating discontinuous outcrop, for more than 100 m (Figs 2 & 3). Horizontal strata mainly consist of thin to very thin beds and laminae (Figs 4C, D & E). Individual strata typically are planar. Sets of horizontal strata may be separated by low angle stratification (Fig. 4C) and they usually grade laterally into concave-upward low angle strata (Site 9 of Fig. 3).

#### Interpretation

Large scale sets of horizontal strata are interpreted to be dune bottom-set deposits and interdune, aeolian sand-sheet deposits formed mainly by grainfall and the movement of wind ripples, plane beds/laminae and dunes across the surface of interdune sand sheets (following Glennie 1970, Hunter 1977, Ahlbrandt & Fryberger 1981, Kocurek 1986, Lancaster 1988). The movement of sand as ripples and plane beds or laminae across sand sheets produces horizontal strata during moderate to strong wind action (Bagnold 1941, Glennie 1970, Jopling 1978). In terms of this study, the direction of sand transport cannot be inferred from horizontal stratification.

Sand flat terrain, if geomorphically analogous to the sand flats of some desert areas (eg McKee & Tibbits 1964, Ahlbrandt & Fryberger 1981), can be expected to be underlain predominantly by horizontal strata, and these strata would pass laterally into dune terrains underlain by cross-strata. That is, cross-stratified dunes typically grade laterally into horizontally stratified sand sheets of interdune areas (Figs 2 & 3). Quarries located in sand hills may be expected to have few horizontal strata. In desert ergs cross-stratified dunes and plinths may pass laterally into horizontally stratified interdunes, and both may be underlain by horizontally stratified sheets of sand.

### Massive structure

#### Description

Sand units with massive structure have no internal stratification. These units occur in three main settings: below the present landsurface to a depth of 3-5 m; as thin sheets typically 0.5-2 m thick beneath buried unconformity surfaces; and up to several metres above the water table (Table 2, Fig. 7). Sequences of sand below the water table have not been observed *in situ*. Units with massive structure commonly exhibit a range of bioturbation features, as will be discussed later.

#### Interpretation

The massive structure of yellow sand is an overprint feature and predominantly the product of bioturbation. Thus massive yellow sand may have originally been stratified. This interpretation is consistent with the yellow sand dunes and sheets being relict, and with the common association of bioturbation overprints with relict desert aeolian deposits (Glennie 1970, Fryberger *et al* 1979, Glassford 1980, 1987; Ward 1988). Buried units of massive yellow sand, which are overlain by stratified yellow sand, indicate an alternating history of desert aeolian deposition followed by bioturbation during subsequent intervening periods of semi-arid or humid conditions. The stratigraphic sequence of structures thereby also is consistent with the concept of alternations from glacial age desert phases to interglacial age humid phases. Such alternations characterise the late Cainozoic of many low and middle latitude areas of the world (Lamb 1977, Sarnthein 1978, Glassford 1980, Thomas & Goudie 1984).

### Major unconformities

#### Description

Major unconformities occur in 13 quarries (Tables 1 & 2; Fig. 7). These unconformities may be within several metres of the present landsurface (Figs 2, 3 & 4), but more commonly occur near or disappear below the floor of quarries. Unconformities have been traced continuously along quarry walls for distances of up to 100 m and by extrapolating discontinuous exposures for over 300 m (Figs 2 & 3). Their observable extent is always limited by the dimensions of the quarry. Deep sequences of yellow sands also have more unconformities than shallow sequences.

Unconformity surfaces are broadly undulose with straight planar segments grading to curved segments (Figs 2, 3 & 4). Straight planar segments may be near-horizontal or inclined at angles of up to 15°. Curved segments are typically concave-up with maximum amplitudes of 5-6m over 50-60m. Unconformities typically have a discordant relationship with more than one set of cross-strata. Furthermore, they also typically separate units of yellow sand which are lithologically distinct in terms of structure (eg cross-stratified or massive), textures (eg coarser or finer, clay-rich or clay poor), composition and colour (eg reddish haematite-pigmented or yellow goethite-pigmented kaolin as coatings on framework grains and as fine sand to silt-clay sized grains). Unconformity surfaces within sequences of yellow sands are commonly underlain by a thin, discontinuous sheet of massive white quartz sand which typically is several centimetres to a few decimetres thick. In places the lower boundary of such sheets has decimetre to metre deep pockets and pipes of white sand, 5-20 cm in diameter, which taper downwards into underlying yellow sands.

**Interpretation**

Disconformities are marked by palaeosols (identified from overprint features; Table 6). They are buried former land surfaces, similar to the present vegetated land surface below which there are pedogenic overprints. These disconformities are super-bounding surfaces (Kocurek 1988), and most are similar to the regional bounding surface (Talbot 1985) category of super-surfaces. Disconform-

ity surfaces have formed by erosion and soil development during interstadial or interglacial intervals under semi-arid or wet conditions similar to the present. Disconformities which are not underlain by a palaeosol, but are erosional, may represent one or more glacial or interglacial intervals. These disconformities are similar to the first order bounding surfaces of Brookfield (1977), and the first and second order bounding surfaces of Kocurek (1981).

**Table 6**

Description and interpretation of secondary alteration features, on or within yellow sand, which have overprinted primary stratification.

<b>Overprint feature</b>	Burrow structure
<b>Description</b>	Occurs up to 3-6m below the surface or below deeply buried surfaces (disconformities); burrow-like structures range from less than 1cm to several centimetres in diameter; depending on the plane of section burrows may be round to cylindrical; burrow fill may differ from surrounding sand in terms of texture, colour and grain types; burrows have a sharp contact with surrounding sand.
<b>Interpretation</b>	Burrowing by various insects, spiders and vertebrates and infilling of burrows when repeated destroys cross-strata and produces a pervasive burrow-mottled structure in detail at the mesoscale, and a massive structure at the macroscale.
<b>Overprint feature</b>	Fibrous root structure (a variety of root structure)
<b>Description</b>	Occurs up to 3-6m below the landsurface or 3-6m below buried disconformities; thin 0.5-1cm diameter vertical to branching cord-like (dendritic) sand, more coherent than surrounding sand when subjected to subaerial exposure; dendritic sand may differ from surrounding yellow sand mainly by being white.
<b>Interpretation</b>	Growth of fibrous plant roots displaces cross-strata and following death and decomposition the root channels are infilled with sand resulting in a dendritic structure of slightly more coherent sand at a mesoscale; at a macroscale the sand appears massive.
<b>Overprint feature</b>	Tap root structure (a variety of root structure)
<b>Description</b>	Occurs to depths of more than 12m below the present landsurface; tap root structures typically are 5-20cm in diameter and this size extends continuously over vertical distances of more than 12m; the structures are typically tube-like with a white sand fill and ferruginous sand rim; the pipe-like structures typically have a vertically discordant relationship with horizontal strata
<b>Interpretation</b>	Growth by tap roots displaces strata; these tap roots act as conduits and sites for solutions and suspensions of iron and aluminium minerals thereby producing multiple vertically discordant pipes of white and ferruginous sand
<b>Overprint feature</b>	Colour mottles
<b>Description</b>	Occur firstly in a broad sheet-like zone up to 6-8m above the groundwater table; secondly in a broad sheet-like zone up to 3-8m below the landsurface or buried landsurfaces; mottles are typically variable in size and shape; mottle diameters range from a few mm to several decimetres, mottle shapes range from equant through pipe-like to highly indented amoeboidal; mottles are typically white quartz sand differing from surrounding yellow sand only in terms of colour and goethite, haematite and kaolin content
<b>Interpretation</b>	Vadose zone and the zone of capillary rise of intermittent wetting and drying and fluctuating Eh and pH in the phreatic zone results in differential pedogenic removal of silt-clay sized coatings of goethite, kaolin and quartz from the surface of sand grains.
<b>Overprint feature</b>	Massive structure
<b>Description</b>	Occurs up to 4-6m below the present landsurface and the surface of buried units of yellow sand, may grade vertically or laterally into units of root structured and/or burrow-structured yellow sand and/or colour mottled sand.
<b>Interpretation</b>	Physical perturbations (eg wetting and drying, removal of fines, settling, compaction) and biological perturbations (eg root-growth, burrowing) destroy cross strata by disrupting and mixing sand layers to form a uniformly homogeneous yellow sand.
<b>Overprint feature</b>	Termitarium (or labyrinthoid) structure
<b>Description</b>	Occurs up to 2-4m below the present landsurface or below buried landsurfaces of yellow or brown sand/sandstone which are relatively rich in silt-clay sized goethite and/or kaolin; labyrinthoid structure comprises a tortuous pattern of vermiform channels to vesicular voids with brown or yellow sand/sandstone (coffee rock) walls and fill; typically weakly coherent; termitarium or labyrinthoid structure typically forms a 0.5 to 1.5m thick sheet roughly parallel to the present landsurface. The upper surface of this sheet typically is c 0.5-2m below the present landsurface.
<b>Interpretation</b>	Termites build a maze of channels with coherent walls, and over time destroy stratification; this results in a sheet of labyrinthoid structured sand/sandstone.

Overprint feature	Brown nodules and brown cementation to form sandstone ( <i>coffee rock</i> )
Description	Occurs mainly in a sheet like zone between 0.5 to 2m thick and tends to be thicker in interdunes; brown to yellowish brown ferruginous quartz sand nodules supported in brown to yellow sand or brown ferruginous, massive, labyrinthoid quartz sandstone, nodules consist of quartz sand supported in a silt-clay matrix of goethite, kaolin and quartz; overlies yellow sand and underlies white sand; upper boundary may be smooth abruptly gradational and broadly parallel with the landsurface or interfingering with white sand over vertical distances of 1-2m, the lower boundary is abruptly gradational to yellow sand and roughly parallel to the upper boundary.
Interpretation	Silt-clay sized matrix material and framework grain coatings of goethite, kaolin and quartz have been removed from yellow sand and transferred in pore water solution and suspension and deposited, in the vadose zone mainly as sandy nodules to produce brown nodular sand, and in the phreatic zone to produce brown sandstone. Both types of overprint are types of <i>lateritic duricrust</i> (locally termed <i>coffee rock</i> ). Those <i>lateritic duricrusts</i> are altered desert-aeolian sediment because they consist of framework grains of desert-aeolian sand surrounded by remobilized and mineralogically partly transformed desert-aeolian dust. The overprints support the view of Playford <i>et al</i> (1975) that some <i>laterites</i> in South-western Australia are of Pleistocene age.
Overprint feature	White quartz sand
Description	White quartz sand is massive, and is without a grain-surface coating of goethite, kaolin and quartz of silt-clay size. White quartz sand occurs in three settings, firstly, it is 1-2m thick where it occurs within a few metres of the landsurface; secondly, it is 0.5m thick where it marks buried former landsurfaces (at discontinuities); thirdly, it occurs near the water table where it is c 1-4m thick. Near the present landsurface white sand thickens from dune tops to dune flanks to interdune flats, it also typically overlies brown nodular sand or brown sandstone and is overlain by humic sand; its upper boundary is smooth and broadly parallel to the landsurface, and gradational into humic sand; its lower boundary is sharp or abruptly gradational, and irregular or deeply (1-2m) penetrative in the form of 10-30cm diameter white-sand filled pipes into underlying brown sand. The upper boundary of white sand in the vicinity of the groundwater table may be sharp or broadly gradational into yellow sand.
Interpretation	Yellow sand has been transformed to white sand by removal of grain coatings of goethite, kaolin and quartz silt-clay in pore water solution and suspension.
Overprint feature	Humic quartz sand
Description	Massive and fibrous-root structured, grey to black quartz sand with plant debris and silt-clay sized organic material; the quartz sand generally is not coated with goethite and kaolin. The unit forms a sheet-like cover 0.1 to 2m thick over white or yellow sand, it increases in thickness from Spearwood Dune ridges and Bassendean Dune tops into flanks and interdune and ridge flats; humic quartz sand overlies white or yellow quartz sand with a gradational contact, its upper boundary is the present landsurface, and its lower boundary is broadly parallel to landsurface.
Interpretation	Yellow or white quartz sand has been impregnated by plant debris and silt-clay sized organic material to form a pedogenic A-horizon.

The occurrence of up to six units of yellow sand separated by five major discontinuities (Table 2, Fig. 7) is interpreted as the record of six major glacial-age desert aeolian phases and five major interglacial-age semi-arid or humid phases in yellow sands of the Perth region. These units are most likely only the upper portion of the record of desert-aeolian sedimentation in SW Australia (see Logan *et al* 1970, Glassford 1980, 1987).

#### Features which overprint sedimentary stratification

Sedimentary stratification in yellow sand grades into, and generally is gradually obliterated by pedogenic and other alteration (or overprint) features (Fig. 10). Overprint features include: burrow structures; root-structures; colour mottles; massive structure; termitarium structures; cementation by brown limonite and clays to form nodules and sandstone; removal of yellow or red grain coatings to form white quartz sand; and humic infiltration/addition to form humic quartz sand (Table 6). The obliteration is gradational. It decreases downwards below the present landsurface and beneath buried discontinuities, and increases towards the groundwater table. Furthermore, thin, deeply buried discontinuity-bounded units of sand lack internal stratification, but may be sharply overlain and gradationally underlain by stratified yellow sands. These relationships indicate that stratification is a primary feature and that it is overprinted by a variety of secondary features. Bioturbation is further described below.

Yellow sand from 3 to 5 m below the present land surface, and sands immediately below the buried surface of a discontinuity, generally are massive to mottled to variably bioturbated. With increasing depth below the pre-


sent land surface and below buried discontinuities, the massive sand eventually grades into stratified sand. Quarry walls expose living tap roots, dead tap roots, sand-filled tap-root-like structures, and sand-filled to open insect burrows that penetrate the layered sands, with sediment-fills infiltrating from higher horizons. Gradational sequences, from laminated sand, through laminated sand with root and burrow structures to mottled sand and massive sand, indicate the progression of overprinting of stratification predominantly by bioturbation (Table 6; Fig. 10). Much of the yellow sand therefore records a history of disruption, mixing, and homogenizing by biological, chemical and physical processes which have overprinted and destroyed primary aeolian stratification.

#### Discussion and conclusions

The results of this study have many and far reaching implications. The pervasive occurrence of sedimentary stratification in thick sequences of yellow sands in the Bassendean and Spearwood dunes has implications for the origin of these sands. This is because the occurrence of stratification supports a primary aeolian origin, and precludes an origin wholly by *in situ* decalcification of limestones.

Dip direction resultants of the cross-strata of yellow sands indicate winds were dominantly from the eastern sector, sub-dominantly from the NW, and to a minor extent from the SW and W. This precludes a wholly coastal-marine derivation for yellow sand because the easterly component is opposite the SW to NE movement of carbonate sands of the modern coastal dunes (Searle & Semeniuk 1985). Presumably the movement of sands that formed Pleistocene coastal dunes (now limestone) also was from





**Figure 10** Summary diagram of the relationships of cross-stratified yellow sand and secondary overprints. **A** Alteration features which overprint and destroy primary cross-stratification. **B** Stages in the development of a multistoried sequence of intercalated cross-stratified sand, massive sand and major disconformities. For simplicity, zones of humification, cementation and bleaching are not shown.

the SW. An easterly wind direction for yellow sand emplacement is consistent with the occurrence of latest Pleistocene to Holocene lunette dunes, of the Swan Coastal Plain, which are located on the western sides of lakes (eg Benger Swamp, Forresdale Lake, White Lake; see Glassford 1980, Beard 1982).

The sequence of stratified units, representing major desert-aeolian phases or ergs, intercalated with the massive units beneath major disconformities suggest that the yellow sands of the Swan Coastal Plain record a mainly middle Pleistocene sequence of desert ergs. During Plio-Pleistocene times there have been more than 20 glacial periods (Kukla 1977, Lamb 1977, Hooghiemstra 1988) during which high latitude areas of the globe were extensively glaciated and sea levels were much lower than at present. The middle to low latitudes (such as the presently humid SW Australia) then were desertic with winds stronger than at present (Lamb 1977, Sarnthein 1978, Glassford 1980, Glennie 1987, Skackleton 1987, Rea & Leinen 1988). Additional desert-erg sequences may be preserved on the Yilgarn Block and possibly under the continental shelf of Western Australia (Glassford 1980, 1987).

Stratified yellow sands, bounded by major disconformities, thus record mainly middle Pleistocene glacial age desert phases, when desert ergs were developed on the exposed continental shelf and the ancestral Swan Coastal Plain. Glacial age desert-erg phases were separated by interglacial age semi-arid to humid phases, with their accompanying development of bioturbated palaeosols on disconformities.

*Acknowledgements* We thank M J Kriewaldt for critically reviewing the manuscript.

## References

- Allen A D 1981 Groundwater resources of the Swan Coastal Plain, near Perth, Western Australia. In: Groundwater resources of the Swan Coastal Plain (ed B R Whelan) CSIRO, 29-80.
- Ahlbrandt T S & Fryberger S G 1981 Sedimentary features and significance of interdune deposits. In: Recent and ancient non-marine depositional environments: models for exploration (ed F G Ethridge & R M Flores) SEPM Spec Publ 31, Tulsa, Oklahoma, 293-314.
- Ash J E & Wasson R J 1983 Vegetation and sand mobility in the Australian desert dunefield. *Zeit Geomorph N F Suppl* 45: 7-25.
- Bagnold R A 1941 *The physics of Blown Sand and Desert Dunes*. Methuen, London.
- Baxter J L 1977 Heavy mineral sand deposits of Western Australia. *Geol Surv W Aust Min Res Bull* 10.
- Beard J S 1982 Late Pleistocene aridity and aeolian landforms in Western Australia. In: Evolution of the flora and fauna of arid Australia (ed W R Barker & P J M Greenslade) Peacock Publ, Adelaide.
- Beard J S 1984 Aeolian landforms, Part 2. In: Beard J S & Springer B S Geographical data from the Vegetation Survey of Western Australia with map of aeolian landforms 1:3,000,000. Occ Paper 2, Vegmap Publ.
- Blakey R C, Peterson F & Kocurek G 1988 Synthesis of late Palaeozoic and Mesozoic eolian deposits of the Western Interior of the United States. *Sediment Geol* 56: 3-125.
- Brindley G W & Brown G (eds) 1980 *Crystal structures of clay minerals and their x-ray identification*. Min Soc London.
- Brookfield M 1970 Dune trends and wind regime in central Australia. *Zeit Geomorph Suppl* 10: 121-153.
- Brookfield M 1977 The origin of bounding surfaces in ancient aeolian sandstones. *Sedimentology* 24: 303-332.
- Dept Conservation & Environment 1980 Atlas of natural resources Darling System Western Australia explanatory text. Univ W Aust Press, Nedlands.
- Folk R L 1974 *Petrology of sedimentary rocks*. Hemphill, Austin, Texas.
- Fryberger S G, Ahlbrandt T S & Andrews S 1979 Origin, sedimentary features and significance of low-angle eolian "sand sheet" deposits, great Sand Dunes National Monument and Vicinity, Colorado. *J Sed Petrol* 49: 733-746.
- Gary M, McAfee R & Wolf C I 1972 *Glossary of geology*. Am Geol Inst Washington D C.
- Glassford D K 1980 Late Cainozoic desert eolian sedimentation in Western Australia. Ph D thesis, Univ W Aust.
- Glassford D K 1987 Cainozoic stratigraphy of the Yeelirrie area, northeastern Yilgarn Block, Western Australia. *J R Soc W Aust* 70: 1-24.
- Glassford D K & Killigrew L P 1976 Evidence for Quaternary westward extension of the Australian Desert into southwestern Australia. *Search* 7: 394-396.
- Glennie K W 1970 *Desert sedimentary environments*. Elsevier, Amsterdam.
- Glennie K W 1987 *Desert sedimentary environments, present and past - a summary*. *Sediment Geol* 50: 135-165.
- Gozzard J R 1983 Fremantle Part Sheets 2033 I and 2033 IV, Perth Metropolitan Region, 1:50,000 Environmental Geology Series, Geol Surv W Aust.
- Gozzard J K 1986 Perth, Sheet 2034 II and part 2034 III and 2134 III, Perth Metropolitan Region, 1:50,000 Environmental Geology Series, Geol Surv W Aust.
- Hooghiemstra H 1988 The orbital-tuned marine oxygen isotope record applied to the middle and late Pleistocene pollen record of Funza (Colombian Andes). *Palaeogeog Palaeoclim Palaeoecol* 66: 9-17.
- Hunter R E 1977 Terminology of cross-stratified sedimentary layers and climbing ripple structures. *Sediment Petrol* 47: 697-706.
- Jopling A V 1978 Bedding genesis. In: *The Encyclopaedia of Sedimentology* (ed R W Fairbridge & J Bourgeois) Dowden, Hutchinson & Ross, Stroudsburg, Penn. 47-56.
- Jordan J E 1986 Armadale, part sheets 2033 I and 2133 IV, Perth Metropolitan Region, 1:50,000 Environmental Geology Series, Geol Surv W Aust.
- Killigrew L P & Glassford D K 1976 Origin and significance of kaolin spherites in sediments of southwestern Australia. *Search* 7: 393-394.
- Klenowski G 1976 Geotechnical properties of the Coastal Limestone in the Perth Metropolitan area. *Geol Surv W Aust Ann Rep* for 1975: 42-46.
- Kocurek G 1981 Significance of interdune deposits and bounding surfaces in aeolian dune sands. *Sedimentology* 28: 753-780.
- Kocurek G 1986 Origins of low-angle stratification in aeolian deposits. In: *Aeolian Geomorphology* (ed W G Nickling), 177-194. Allen & Unwin, London.
- Kocurek G 1988 First-order and super bounding surfaces in eolian sequences - Bounding surfaces revisited. *Sediment Geol* 56: 193-206.
- Kukla G J 1977 Pleistocene land-sea correlations, I. Europe. *Earth Sci Rev* 13: 307-374.
- Lamb H H 1977 *Climate present, past and future, volume 2, climatic history and the future*. Methuen, London.
- Lancaster N 1988 The development of large aeolian bedforms. *Sediment Geol* 55: 69-89.
- Leeder M R 1982 *Sedimentology*. Allen & Unwin, London.
- Lindholm R C 1987 *A practical approach to sedimentology*. Allen & Unwin, London.
- Logan B W, Read J F & Davies G R 1970 History of carbonate sedimentation, Quaternary epoch, Shark Bay, Western Australia. *Am Assoc Petrol Geol Mem* 13: 38-84.
- Low G H 1971 Definition of two Quaternary formations in the Perth Basin. *Geol Surv W Aust Ann Rep* for 1970: 33-34.
- Low G H & Lake R W 1970 Perth and environs geological maps, sheet 2: 1:50,000 scale. *Geol Surv W Aust*.
- Low G H, Lake R W, Baxter J L & Doepel J J C 1970 Perth and environs geological maps, sheet 3: 1:50,000 scale. *Geol Surv W Aust*.
- Lowry D C 1977 Perth Basin yellow sand. *Search* 8: 54-55.
- Marzoff J E 1988 Controls on late Paleozoic and early Mesozoic eolian deposition of the Western United States. *Sediment Geol* 56: 167-191.
- McArthur W M & Bettenay E 1960 The development and distribution of the soils of the Swan Coastal Plain, Western Australia. CSIRO Soil Publ 16.
- McKee E D & Bigarella J J 1979 Sedimentary structures in dunes with two sections on the Lagoa dune field, Brazil, 83-136. In: *Geol Surv Prof Paper* 1052, US Gov Printing Office, Washington.
- McKee E D & Tibbitts 1964 Primary structures of a self dune and associated deposits in Libya. *J Sediment Petrol* 34: 5-17.
- McKee E D & Weir G W 1953 Terminology for stratification and cross-stratification in sedimentary rocks. *Geol Soc Am Bull* 64: 381-390.
- Nemec W 1988 The shape of the rose. *Sediment Geol* 59: 149-152.
- Nielson J & Kocurek G 1987 Surface processes, deposits and development of star dunes: Dument dune field, California. *Geol Soc Am Bull* 99: 177-186.
- Passmore J R 1970 Shallow coastal aquifers in the Rockingham district, Western Australia. *Water Res Found Aust Bull* 18: 83p.
- Pettijohn F J, Potter P E & Siever R 1987 *Sand and sandstone*, 2nd ed. Springer-Verlag, New York.
- Playford P E, Cockbain A E & Low G H 1976 Geology of the Perth Basin Western Australia. *Geol Surv W Aust Bull* 124.
- Playford P E, Cope R N, Cockbain A E, Low G H & Lowry D C 1975 Phanerozoic in Geology of Western Australia. *W Aust Geol Surv Mem* 2: 223-433.
- Playford P E & Low G H 1972 Definitions of some new and revised rock units in the Perth Basin. *Geol Surv W Aust Ann Rep* for 1971: 44-46.